

Suggestions on TAP

LIMSEOK CHANG

National Institute of Environmental Research, Korea

TAP, 9~ 10 July, Russia

1. Background
2. Suggestions
3. Discussion



1. Background and Objectives

No Comprehensive mechanism in NE Asia

Targets	NE Asia	Europe
1. Acid deposition, Eeutrophication	○(EANET, LTP)	EMEP
2.Photochemical Oxidnats	○ (LTP, TEMM Project)	EMEP
3.Heavy metals	○(LTP)	EMEP
4.POPs	△(East Asia POPs Monitoring)	EMEP
5.PM	○(LTP)	EMEP
6. Integrated Modeling	×	EMEP
7.Emission Inventory	△(LTP, MICS-ASIA)	EMEP
8.Emission Process Model	○(LTP)	EMEP
9.Openness	×	EMEP
10.Collboration with HF-TAP	×	EMEP (HTAP)

LTP Background

- LTP was started in year 2000 as a **government-based air pollution research framework** among China, Japan, and Korea
- LTP's activities have been mostly focusing on understanding **transboundary air pollution** and **S-R relationship among three countries**, using modeling and monitoring techniques
- Now LTP members are planning the **4th year of the 3rd Stage (2012)**



Similarities and uniqueness of LTP Framework to other activities


- **Similarity**

- Air Quality Monitoring for Asia (to EANET)
- Air Quality S-R Modeling for Asia (to MICS-Asia)
- Target pollutants – Sulfur, Nitrogen, and others (EANET and MICS-Asia)
- RAINS-Asia, GAINS-Asia, ABC, Global-Chem Modeling, and others...

- **Uniqueness**

- Government-initiated scientific research collaboration framework in support of regional air quality issues
- Both modeling and monitoring
- Both pure science and policy supporting science
- Strictly focus on East Asia (three countries)
- Long lasting geo-scientific collaboration in East Asia

Model weakness (Model domain and locations of measurement)



• six SO₂ monitoring sites, and aircraft pathways







LTP Modeling framework

	China	Japan	Korea
Model system	Models-3 / CMAQ σ coordinate 14 layers, 70×66 grids, 60km resolution (Byun and Ching, 1999)	RAQM (Regional Air quality Model) terrain following coordinate 12 layers, 110×80 grids, 60km resolution (An et al., 2002)	CADM (Comprehensive Acid Deposition Model) terrain following coordinates 12 layers, 110×80 grids, 60km resolution (Lee et al., 1998)
Domain	20 ~ 50N, 100 ~ 150E	20 ~ 50N, 100 ~ 150E	20 ~ 50N, 100 ~ 150E
Meteorological Model	MM5 34 layers with FDDA using NCEP reanalysis	MM5 125×95 (45km), 23 layers, FDDA using NCEP FNL reanalysis	CSU-RAMS 110×80, 29 vertical layer FDDA using NCEP FNL reanalysis
Chemical Mechanism	RADM Chemistry	CBM-IV mechanism	RADM Chemistry
Cloud Model Physical option	Diagnostic cloud model in RADM Simple explicit moisture scheme Grell cumulus schemes, MRF	Cloud model in MM5 Betts-Miller cumulus scheme, MRF RRTM	Cloud model in CSU-RAMS Anthes-Kuo cumulus scheme, MRF
Emission	SO ₂ , NO _x , VOC, NH ₃ , CO, PM ₁₀ , biogenic VOC provided by LTP for the base year of 1998 (1°×1° resolution)	Same as China	Same as China
Dry deposition	Wesely's parameterization (Wesely, 1989)	Modified Wesely's parameterization (Walmsley & Wesely, 1996)	Dry deposition module in RADM (Lee et al, 1998)
Wet deposition	RADM Module (Chang et al, 1987)	RADM Module (Chang et al, 1987)	RADM Module (Chang et al, 1987)
Land use type	EPA/NOAA global ecosystem (11 categories)	DeFries & Townshend (1994)	EPA/NOAA global ecosystem (11 categories)


Summary of model performance (SO2)

	Obs	China	Japan	Korea
Sample size	57	57	57	57
Range(ppb)	0.0042-35.43	0.0-11.387	0.05-16.07	0.025-24.22
Mean(ppb)	4.59	2.35	2.91	5.01
Standard deviation (ppb)	7.13	2.66	3.26	5.81
Mean of ratio model/obs (S/O)		0.91	1.99	4.25
Standard deviation of ratio model/obs (S/O)		1.37	6.81	19
Absolute gross error		3.11	3.61	3.44
Correlation coefficient		0.54	0.19	0.53
Mean difference		2.24	1.68	-0.42
Difference standard deviation		6.13	7.24	6.42
Root-mean square error		6.48	7.37	6.38
Mean square error. MSEN		5.06	10.21	24.56
Mean square error. MSES		37.54	45.05	16.86
Index of agreement		0.53	0.37	0.69
Mean fractional error		0.59	0.22	-0.12

Model performance comparison



Model performance comparison




Spatial distribution of the simulated [SO₂]

a) CHINA


b) JAPAN

c) KOREA




Simulated surface SO₂ concentrations from (a) China, (b) Japan, and (c) Korea on 5-13 March.

Conversion ratio of SO₂ to sulfate




Time variations of longitudinal cross-section of simulated conversion ratio of sulfur ($F_s = \text{SO}_4^{2-}/(\text{SO}_2 + \text{SO}_4^{2-})$) from (a) China, (b) Japan, and (c) Korea on 5-13, March.

Spatial distribution of the simulated [SO₂]



Seasonally averaged surface SO₂ and SO₄²⁻ concentrations simulated by (a) China, (b) Japan, and (c) Korea for the year of 2002.

Model validation by observations




Comparison between monthly variation of simulated versus observed SO₂ concentrations at 6 LTP sites in 2002

New Challenges

- The **LTP research framework** has been **working out**, but the **data and methodology for research are outdated** from scientific viewpoint and **research components are not complete** in policy supporting viewpoint

• Climate change




- climate influence on air quality
- future scenario
- co-benefits

• Local and regional air pollution



- Inter- vs. intra- national S-R
- Megacity impacts

• Category Integration



- CAPs + GHGs + HAPs
- Impact study



• Understand pollution events



- Dust storm, wildfire, storms, heat
- Air quality forecasting

Trend

Night light during last 10 years




Rapid economic growth

Transboundary Air Pollution Problem


→ Collaboration among stakeholders

Population and Urbanization


ROK



DPRK




China




Target year?


Japan



Urbanization



Time and Emissions



Source : NIER, 2011

New Objectives

- Two major and one supplemental objectives

: Understand **air quality issues** in East Asia in consideration of **new challenges**, such as **secondary pollutants, HAPs, climate change**, and etc. Decide what we want **to pursue** and what we **won't** (**State-of-art science**)

: Use our understanding **to prioritize our actions to mitigate** adverse AQ effects for another decade. Health/environmental **impact** and **mitigation policy** study need to be initiated (**Policy supporting science**)

: How can we **improve our collaborative research framework** to accomplish these objectives effectively?

2. Suggested Topics

- Air Quality Forecast for Northeast Asia
- Implementation of Advanced S-R Methodologies
- Assessment of O₃ and PM
- Scenario-based Collaboration Simulation Approach

2.1 Chemical Air Quality Forecast for Northeast Asia

Air Quality Forecast - Existing Efforts (examples)

Modeling Frameworks

- Meteorological Models : MM5, RAMS, WRF
- Chemical Models : CMAQ, CFORS, CAMx
- Domain, Grid, and Emissions

✓ China



- Horizontal Grid Spacing:
Asia: 36km, China and Korea:
12km, Beijing Metropolitan Area: 4km
- Vertical Layer : 14 layers
- Emissions : Trace-P 2000 + Statistical Books

✓ Japan

- Horizontal Grid Spacing:
Asia: 80km , 15 km (in preparation)
- Vertical Layer : 23 layers to 20km
- Emissions : Trace-P 2000 + dust, sea salt, lightning, volcano, radon, Biomass burning

✓ Korea

- Horizontal Grid Spacing:
East Asia: 27km , Korea: 9km, SMA: 3 km
- Vertical Layer : 11 layers to 14.7km
- Emissions : INTEX-B 2006+ CAPSS 2007+ fugitive dust, biogenic, biomass burning



Air Quality Forecast – a Framework (Draft)

- Three country generate the common metrological/emissions data for the simulation
- Each country uses its' own chemical model(s)
- Cooperation with other fields (ground and airborne monitoring, satellite data retrieval, LIDAR networks)



- **Integrated performance testing during Intensive Monitoring Period (ex. forecast for 72 hours for the entire month of IMP)**
 - National Air Quality Index (API)
 - Concentration of air pollutants : O_3 , PM_{10} , $PM_{2.5}$, SO_2 , NO_2 , sulfate, nitrate, ammonia, mercury etc.



Need common /integrated modeling/monitoring framework and may need a modeling center

For performance enhancement

- Input data update (Emission Inventory, Land Use, Terrain)
- More detailed emission Inventory (Fugitive dust, Sand storm, Forest fire, Biomass burning, Volcano, Biogenic emission)
- Emission inventory for North Korea
- Background, Boundary Conditions

For implementation - Arrangements

- Each country secures its' financial support from Government and/or International Organizations.
- Each country reports to the Environmental Minister and endeavors to bring this matter to the Tripartite Environment Ministers Meeting Among China, Japan, and Korea (TEMM)

2.2 Implementation of Advanced S-R Methodologies

Why Do We Need Advanced S-R Methodologies?


- **To conduct more sophisticate S-R research**
 - Simultaneous analysis for regions-sectors, ICs, BCs
 - Finer S-R(e.g. megacity impact study)
 - Control measure testing ...
- **To understand various sensitivities**
 - Precursor to product sensitivity
 - Region to region sensitivities...
- **Benefits**
 - Save time, effort, and space(faster and simpler)
 - Avoid non-linearity

What's available?

- Source tagging(REMSAD, CAMx-PSAT, CMAQ-PPTM)
- Forward Sensitivity(CMAQ-DDM)
- and more...


Source Tagging(CMAQ-PPTM)

- PPTM : Particle and Precursor Tagging Methodology
- Assessment of the source contribution by **source tagging** method
- Emissions from selected sources, source categories, or source regions are (numerically) tagged and then tracked throughout a simulation



Source Tagging(REMSAD-tagging)

Woo et al.(2006)



Assessment of megacity impacts

2.3 Assessment of O₃ and PM for the future LTP Project

Key Findings of O₃ Impacts (from TF-HTAP 2010)

(TF-HTAP :Task Force on Hemispheric Transport of Air Pollution)

O₃ Impacts on Health Impacts

- One study based on the HTAP multi-model comparison estimated that O₃ resulting from emissions from foreign regions contributes 20% to >50% of O₃ mortalities, subject to large uncertainty
- Three studies estimate that reductions in O₃ precursor emissions may avoid more premature mortalities outside of some source regions than within, mainly because of larger populations outside of the source regions

O₃ Impacts on Climate Change

- O₃ contributes significantly to climate forcing,
 - Directly as a greenhouse gas that causes warming
 - and indirectly by damaging plants inhibiting their natural uptake of CO₂.
- Among ozone precursors, widespread reductions in emissions of CH₄, CO, and VOCs better reduce net climate forcing than reducing NO_x, which may increase forcing over decadal time scales.

Key Findings of PM Impacts (from HTAP 2010)

PM Impacts on Health

- Contributions to PM from emissions within a region are much more important for health than emissions from foreign continents
- Intercontinental transport of PM can cause more mortalities than intercontinental transport of O₃, due to the stronger PM-mortality relationship.
- Emissions from North America and Europe have much greater impacts on foreign regions than do emissions from East Asia and South Asia

Hundreds of annual avoided mortalities, threshold=5.8 µg/m³ in *italics*

Source Region	Receptor Region				
	NA	EA	SA	EU	World
NA	502	20	9	49	590
	<i>125</i>	<i>19</i>	<i>8</i>	<i>38</i>	<i>190</i>
EA	10	4348	25	18	4433
	<i>5</i>	<i>3330</i>	<i>23</i>	<i>14</i>	<i>3376</i>
SA	2	42	2105	5	2168
	<i>1</i>	<i>39</i>	<i>1099</i>	<i>3</i>	<i>1142</i>
EU	8	82	70	1769	2010
	<i>4</i>	<i>78</i>	<i>57</i>	<i>573</i>	<i>71600</i>

Impact on foreign receptor regions:

← 14.9%

← 1.9%

← 2.9%

← 12.0%



Key Findings of PM Impacts (from HTAP 2010)

PM Impacts on Climate Change

- Change of global annual average TOA all-sky aerosol direct RF in response to the 20% reduction of anthropogenic emissions
- BC activity under EMEP/CLRTAP: Focus country specific contribution to direct radiative forcing by BC aerosols

(unit: mW m⁻², mean ± std. dev)

Source Region	Sulfate	POM	BC	Sulfate+POM+BC
NA	16.1 ±5.6	1.6 ±1.0	-4.5 ±1.9	13.2 ±5.2
EU	26.7 ±9.5	1.9 ±1.2	-7.4 ±2.3	21.2 ±9.5
EA	19.6 ±7.2	3.2 ±1.8	-14.5 ±8.0	8.4 ±10.2
SA	6.1 ±1.9	2.5 ±1.3	-5.5 ±2.4	3.1 ±3.2
NA+EU+EA+SA	68.4 ±22.9	9.1 ±5.0	-31.9 ±13.7	45.9 ±24.6

2.4 Scenario-based Collaboration Simulation Approach

Roadblocks for international environmental cooperation

▪ Among countries

- Lack of trust, data sharing
- Different (political and economic) interests
- Lack of legitimacy of independent modeling

▪ Among decisionmakers and scientists

- Lack of understanding
- Different assumption, languages and interests

Conditions for effective collaboration

- **Existence of on-going** (effective and flexible) **communication channels** for domestic and international decisionmakers, key stakeholders, and scientists
- **Joint Fact-finding mechanism**
 - Shared assumptions in modelling
 - Shared data and research methods
- **Shared roadmap** from the beginning

Alternative approach to conventional ways

▪ Conventional channels

- Government-initiated (established) meetings
- Scientists-oriented meetings

▪ Alternative channels

- **Workshop environment** where decisionmakers, key stakeholders, and scientists **get together**
- More **flexible and creative** environment
- For learning about the relationships between **science and policies and politics**

Scenario-based simulation method

- **Construction of potential scenarios regarding transboundary air pollution** (and/or) climate change with scenario development team among three countries (China, Japan, and Korea)
- **Integration of Scenario with Modeling**
- Utilization of scenario-based modeling with **collaboration simulation exercise**

Potential benefits of alternative approach

- **Lower the tension** among participants
- Promote comprehensive understanding about **complex relationships on the issue**
- **Focus on future** rather than the past
- **Facilitate understanding of non-scientists** on the model and its outcome
- **Improve model design** with inputs from other stakeholders and decisionmakers
- Maintain the communication among various actors
- Bridge the **flexible communications** with formal ones

Pre-requisite

- Acknowledgement of the potential benefits by key decision makers in three countries
- Identification of neutral convener or facilitators, experts on scenario-based planning in each country
- Knowledge on scientific, economic, political, policy factors

Thank you for Attention!
